



University of Belgrade, Technical Faculty in Bor
29th International Conference Ecological Truth
& Environmental Research



EcoTER'22

Proceedings



Editor

Prof. Dr Snežana Šerbula

21-24 June 2022, Hotel Sunce, Sokobanja, Serbia



University of Belgrade, Technical Faculty in Bor
29th International Conference Ecological Truth
& Environmental Research



EcoTER'22

Proceedings



Editor

Prof. Dr Snežana Šerbula

21-24 June 2022, Hotel Sunce, Sokobanja, Serbia

PROCEEDINGS

29th INTERNATIONAL CONFERENCE

ECOLOGICAL TRUTH AND ENVIRONMENTAL RESEARCH – EcoTER'22

Editor:

Prof. Dr Snežana Šerbula

University of Belgrade, Technical Faculty in Bor

Technical Editors:

Dr Jelena Milosavljević

University of Belgrade, Technical Faculty in Bor

Asst. Prof. Dr Ana Radojević

University of Belgrade, Technical Faculty in Bor

Dr Jelena Kalinović

University of Belgrade, Technical Faculty in Bor

Asst. Prof. Dr Tanja Kalinović

University of Belgrade, Technical Faculty in Bor

Asst. Prof. Dr Žaklina Tasić

University of Belgrade, Technical Faculty in Bor

Publisher: University of Belgrade, Technical Faculty in Bor

For the Publisher: Prof. Dr Nada Štrbac, Dean

Printed: GRAFIK CENTAR DOO Beograd, 120 copies

Year of publication: 2022

ISBN 978-86-6305-123-2

CIP - Каталогizacija u publikaciji
Narodna biblioteka Srbije, Beograd

502/504(082)(0.034.2)

574(082)(0.034.2)

INTERNATIONAL Conference Ecological Truth & Environmental Research (29 ; 2022 ; Sokobanja)

Proceedings [Elektronski izvor] / 29th International Conference Ecological Truth and Environmental Research - EcoTER'22, 21-24 June 2022, Sokobanja, Serbia ; [organized by University of Belgrade, Technical faculty in Bor (Serbia)] ; [co-organizers University of Banja Luka, Faculty of Technology – Banja Luka (B&H) ... [et al.]] ; editor Snežana Šerbula. - Bor : University of Belgrade, Technical faculty, 2022 (Beograd : Grafik centar). - 1 USB fleš memorija ; 5 x 5 x 1 cm

Sistemski zahtevi: Nisu navedeni. - Nasl. sa naslovne strane dokumenta. - Tiraž 120. - Bibliografija uz svaki rad. - Registar.

ISBN 978-86-6305-123-2

a) Животна средина -- Зборници б) Екологија -- Зборници

COBISS.SR-ID 69053705

**29th International Conference
Ecological Truth and Environmental Research 2022**

is organized by:

**UNIVERSITY OF BELGRADE, TECHNICAL FACULTY IN
BOR (SERBIA)**

Co-organizers of the Conference:

**University of Banja Luka, Faculty of Technology
– Banja Luka (B&H)**

**University of Montenegro, Faculty of Metallurgy and Technology
– Podgorica (Montenegro)**

**University of Zagreb, Faculty of Metallurgy
– Sisak (Croatia)**

**University of Pristina, Faculty of Technical Sciences
– Kosovska Mitrovica**

Association of Young Researchers – Bor (Serbia)



**The Conference is financially supported by
the Ministry of Education, Science and Technological
Development of the Republic of Serbia**

Platinum Donor of the Conference



塞尔维亚紫金铜业有限公司
SERBIA ZIJIN COPPER DOO

HONORARY COMMITTEE

Dr Petar Paunović, Zaječar (Serbia)
Prof. Dr Stevan Stanković, Bor (Serbia)
Prof. Dr Zvonimir Stanković, Bor (Serbia)
Prof. Dr Velizar Stanković, Bor (Serbia)
Dragan Randelović, Society of Young Researchers – Bor (Serbia)
Toplica Marjanović, Society of Young Researchers – Bor (Serbia)
Mihajlo Stanković, Special Nature Reservation of Zasavica, Sremska Mitrovica (Serbia)

SCIENTIFIC COMMITTEE

Prof. Dr Snežana Šerbula, <i>President</i>	
Prof. Dr Alok Mittal (India)	Prof. Dr Yeomin Yoon (United States of America)
Prof. Dr Jan Bogaert (Belgium)	Prof. Dr Chang-min Park (South Korea)
Prof. Dr Aleksandra Nadgórska-Socha (Poland)	Prof. Dr Faramarz Doulati Ardejani (Iran)
Prof. Dr Luis A. Cisternas (Chile)	Prof. Dr Ladislav Lazić (Croatia)
Prof. Dr Wenhong Fan (China)	Prof. Dr Natalija Dolić (Croatia)
Prof. Dr Martin Brtnický (Czech Republic)	Prof. Dr Milutin Milosavljević (Kosovska Mitrovica)
Prof. Dr Isabel Maria De Oliveira Abrantes (Portugal)	Prof. Dr Nenad Stavretović (Serbia)
Prof. Dr Shengguo Xue (China)	Prof. Dr Ivan Mihajlović (Serbia)
Prof. Dr Tomáš Lošák (Czech Republic)	Prof. Dr Milovan Vuković (Serbia)
Prof. Dr Maurice Millet (France)	Prof. Dr Nada Blagojević (Montenegro)
Prof. Dr Murray T. Brown (New Zealand)	Prof. Dr Darko Vuksanović (Montenegro)
Prof. Dr Xiaosan Luo (China)	Prof. Dr Irena Nikolić (Montenegro)
Prof. Dr Daniel J. Bain (United States of America)	Prof. Dr Šefket Goletić (B&H)
Prof. Dr Che Fauziah Binti Ishak (Malaysia)	Prof. Dr Džafer Dautbegović (B&H)
Prof. Dr Richard Thornton Baker (United Kingdom)	Prof. Dr Borislav Malinović (B&H)
Prof. Dr Mohamed Damak (Tunisia)	Prof. Dr Slavica Sladojević (B&H)
Prof. Dr Jyoti Mittal (India)	Prof. Dr Nada Šumatić (B&H)
Prof. Dr Miriam Balaban (United States of America)	Prof. Dr Mirjana Rajčić Vujasinović (Serbia)
Prof. Dr Fernando Carrillo-Navarrete (Spain)	Prof. Dr Snežana Milić (Serbia)
Prof. Dr Pablo L. Higuera (Spain)	Prof. Dr Dejan Tanikić (Serbia)
Prof. Dr Mustafa Cetin (Turkey)	Prof. Dr Milan Trumić (Serbia)
Prof. Dr Mauro Masiol (Italy)	Dr Jasmina Stevanović (Serbia)
Prof. Dr George Z. Kyzas (Greece)	Dr Dragana Đorđević (Serbia)
Prof. Dr Mustafa Imamoğlu (Turkey)	Dr Jelena Milojković (Serbia)
Prof. Dr Petr Solzhenkin (Russia)	

ORGANIZING COMMITTEE

Prof. Dr Snežana Šerbula, *President*

Prof. Dr Snežana Milić, *Vice President*

Prof. Dr Đorđe Nikolić, *Vice President*

Prof. Dr Milica Veličković (Serbia)

Prof. Dr Danijela Voza (Serbia)

Prof. Dr Maja Nujkić (Serbia)

Dr Ana Simonović (Serbia)

Dr Tanja Kalinović (Serbia)

Dr Ana Radojević (Serbia)

Dr Žaklina Tasić (Serbia)

Dr Jelena Kalinović (Serbia)

Dr Jelena Milosavljević (Serbia)

Dr Dragana Medić (Serbia)

Aleksandra Papludis, MSc (Serbia)

Sonja Stanković, MSc (Serbia)

Vladan Nedelkovski, MSc (Serbia)

Mara Manzalović, English language teacher (Serbia)

Enisa Nikolić, English language teacher (Serbia)

PREFACE

In today's world, the environment has been endangered by the use of outdated technology, fossil fuels and environmental law violations. Therefore, environmental and many other scientists all over the world have been concerned about finding sustainable technology in resolving these issues. That is why environmental research and ecological truth are at the focus of the 29th International Conference Ecological Truth & Environmental Research 2022 (EcoTER'22), which will be held in Sokobanja, Serbia, 21–24 June 2022. On behalf of the Organizing Committee, it is a great honor and pleasure to wish all the participants a warm welcome to the Conference.

We hope to convey the message of the conference, which is that a transformation of attitudes and behavior would bring the necessary changes. This is also an opportunity for the participants who are experts in this field to exchange their experiences, expertise and ideas, and also to consider the possibilities for their collaborative research.

The 29th International Conference Ecological Truth & Environmental Research 2022 is organized by the University of Belgrade, Technical Faculty in Bor, and co-organized by the University of Banja Luka, Faculty of Technology, the University of Montenegro, Faculty of Metallurgy and Technology – Podgorica, the University of Zagreb, Faculty of Metallurgy – Sisak, the University of Pristina, Faculty of Technical Sciences – Kosovska Mitrovica and the Association of Young Researchers, Bor.

These proceedings include 85 papers from the authors coming from the universities, research institutes and industries in 6 countries: Bulgaria, Italia, Albania, Bosnia and Herzegovina, Montenegro and Serbia.

As a part of this year's conference, the 4th Student section – EcoTERS'22 is being held. We appreciate the contribution of the students and their mentors who have also participated in the Conference.

Financial assistance provided by the Ministry of Education, Science and Technological Development of the Republic of Serbia is gratefully acknowledged by the Organizing Committee of the EcoTER'22 conference.

The support of the Platinum donor and their willingness and ability to cooperate have been of great importance for the success of EcoTER'22. The Organizing Committee would like to extend their appreciation and gratitude to the Platinum donor of the Conference for their donation and support.

We appreciate the effort of all the authors who have contributed to these Proceedings. We would also like to express our gratitude to the members of the scientific and organizing committees, reviewers, speakers, chairpersons and all the Conference participants for their support to EcoTER'22. Sincere thanks go to all the people who have contributed to the successful organization of EcoTER'22.

Prof. Snežana Šerbula,

President of the Organizing Committee

TABLE OF CONTENTS

Plenary Lectures

<i>Marija Petrović Mihajlović, M. Antonijević</i> PURINES AS GREEN CORROSION INHIBITORS	3
<i>Milan Radovanović, M. Antonijević</i> ENVIRONMENTALLY SAFE CORROSION INHIBITORS: AMINO ACIDS	12
<i>Hyusein Yemendzhiev, Y. Mersinkova, R. Koleva, G. Peeva, V. Nenov</i> BIO-ELECTROCHEMICAL SYSTEMS – STATE OF THE ART: BIOLOGY, ELECTROCHEMISTRY AND APPLICATION IN WASTEWATER MANAGEMENT	25
<i>Aleksandar Marinković</i> GREEN ORGANIC CHEMISTRY: A FRAMEWORK FOR SUSTAINABLE ENVIRONMENTAL PROTECTION	32

Conference Papers

<i>Sanja Mrazovac Kurilić, A. Ćirišan, Z. Podražčanin, Lj. Nikolić Bujanović</i> SO ₂ POLLUTION IN ŠABAC (2009–2020)	39
<i>Ana Čučulović, J. Stanojković, R. Čučulović, S. Nestorović, N. Radaković</i> THE DISTRIBUTION OF THE MASS CONCENTRATIONS OF K, Th AND U IN THE SOILS OF THE TEKIJIA REGION, THE NP ĐERDAP	43
<i>Jovana Bošnjaković, M. Bugarčić, M. Milošević, N. Prlainović, R. Salih, P. Batinić, A. Popović, M. Đolić</i> APPLICATION OF NANO- MnO ₂ MODIFIED LIGNIN – BASED ADSORBENT FOR REMOVAL OF DICHROMATE IONS AND DICLOFENAC FROM WATER	49
<i>Branislava Matić, S. Živković Perišić, D. Jovanović, S. Dejanović, D. Miljuš, L. Kukobat</i> MAPPING HEALTH RISKS OF CRITERIA CHEMICAL(S) TO DEMONSTRATE BENEFITS OF RISK ASSESSMENT FOR DECISION- MAKERS	55
<i>Violeta Babić, B. Kanjevac, M. Milenković, S. Stajić, M. Vukin, N. Stavretović, M. Račić</i> CHARACTERISTICS OF WINTER TEMPERATURE REGIME IN SPRUCE FOREST ON KOPAONIK	62
<i>Dragana Pavlović, D. Čakmak, V. Perović, M. Matić, M. Marković, M. Mitrović, P. Pavlović</i> ENVIRONMENTAL RISK ASSESSMENT OF PTES IN AGRICULTURAL SOILS AFFECTED BY INDUSTRIAL ACTIVITIES IN BELGRADE	68
<i>Olga Kostić, G. Gajić, S. Jarić, Z. Mataruga, D. Sekulić, N. Radulović, M. Mitrović, P. Pavlović</i> ANALYSIS OF As AND Pb ACCUMULATION IN GARDEN SOIL AND VEGETABLE CROPS IN THREE BELGRADE MUNICIPALITIES	75

Marija Ječmenica Dučić, D. Vasić Aničijević, B. Savić, D. Aćimović, M. Simić, D. Maksin, T. Brdarić	NEW STRATEGIES FOR DEVELOPMENT OF HIGHLY SELECTIVE MATERIALS FOR CARBON DIOXIDE CAPTURE	81
Miljan Bigović, D. Đurović, I. Nikolić, Lj. Ivanović, B. Bajić	CHARACTERISTIC, SOURCE AND ECOLOGICAL RISK OF PAHs IN AGRICULTURAL SOILS PLJEVLJA MUNICIPALITY (MONTENEGRO)	85
Jelena Vranković, K. Jovičić, V. Đikanović	EFFECT OF DIFFERENT ENVIRONMENTAL CONDITIONS ON LIPID PEROXIDATION LEVEL IN <i>Rutilus rutilus</i> (ACTINOPTERYGII: CYPRINIDAE)	91
Aurora Bakaj (Čizmja), M. Lika (Čekani)	EVALUATION OF MICROBIAL ENVIRONMENT ON THE BEACH SAND	96
Dragana Randelović, R. Pantović	IMPACT OF THE MINING ACTIVITIES IN THE GOLIJIA NATURE PARK AREAS SITUATED ON THE TERRITORY OF RAŠKA MUNICIPALITY	104
Bojana Tubić, K. Zorić, N. Popović, M. Raković, N. Marinković, M. Paunović	INDICATIVE ECOLOGICAL STATUS ASSESSMENT OF SELECTED STREAMS ON ROGOZNA MOUNTAIN BASED ON AQUATIC MACROINVERTEBRATES	110
Božica Vasiljević, J. Tomović, A. Atanacković, R. Petrović	INDICATIVE ECOLOGICAL STATUS ASSESSMENT BASED ON EPILOTHIC DIATOMS OF SMALL RIVERS AT ROGOZNA MOUNTAIN	116
Đuro Čokeša, M. Marković, N. Potkonjak, B. Kaluđerović, S. Radmanović, S. Šerbula	ARSENITE-SOIL HUMIC ACID BINDING BY ISOTHERMAL TITRATION CALORIMETRY: THERMODYNAMICS AND MNIS MODEL	121
Tatjana Anđelković, I. Kostić Kokić, B. Zlatković, D. Anđelković	Cu(II) ACCUMULATION POTENTIAL OF AQUATIC MACROPHYTE <i>PISTIA STRATIOTES</i>	127
Tatjana Anđelković, D. Bogdanović, I. Kostić Kokić, H. Kocić, G. Kocić	PHTHALATES MIGRATION FROM ABSORBABLE SURGICAL SUTURES INTO MODEL SOLUTION	132
Aleksandra Papludis, S. Alagić, S. Milić, I. Zlatanović, M. Filipović, J. Nikolić, V. Stankov Jovanović	THE CONTENT OF DANGEROUS CONTAMINANTS PAHs IN THE SOIL AND ROOTS OF <i>HEDERA HELIX</i> IN SLATINA (BOR'S MUNICIPALITY)	137
Jelena Milosavljević, S. Šerbula, A. Radojević, T. Kalinović, J. Kalinović, B. Spalović	TOXIC METALS BIOACCUMULATION IN <i>Plantago lanceolata</i> FROM ANTHROPOGENICALLY DISRUPTED AREA	142
Mirjana Ocokoljić, Dj. Petrov, N. Galečić, M. Miodrag	ECOLOGICAL AND AESTHETIC CHARACTERISTICS OF TREES IN "BELGRADE WATERFRONT" PARKING LOTS	149
Mirjana Ocokoljić, Dj. Petrov, A. Tutundžić, D. Skočajić, S. Petrović	STATE, FUNCTIONS AND PERSPECTIVES OF DENDROFLORA IN THE SCHOOLYARD OF THE MEDICAL SCHOOL IN BELGRADE	154

Mirjana Ocokoljić, Dj. Petrov TRACHYCARPUS FORTUNEI (HOOK.) H. WENDL. IN THE CONDITIONS OF CHANGED TEMPERATE CONTINENTAL CLIMATE IN BELGRADE	159
Marija Perkunić, S. Vukić, Z. Živković, V. Stupar, M. Saulić GREEN OASIS EFFECT ON URBAN ENVIRONMENT	164
Anđela Stojić, D. Tanikić APPLICATION OF GREEN AREAS AND GREEN ROOFS IN URBAN AREAS	169
Gordana Šekularac, T. Ratknić, M. Aksić, N. Gudžić, M. Vranešević, M. Ratknić SOIL EROSION ASSESSMENT USING EPM: A CASE STUDY IN THE CATCHMENT AREA OF THE TURKISH BROOK, WESTERN SERBIA	174
Nataša Knežević, A. Jovanović, R. Salih, Z. Veličković, A. Popović, P. Batinić, A. Marinković, J. Gržetić MODIFIED LIGNIN-BASED MICROSPHERES AS A GREEN SORBENT FOR THE REMOVAL OF CHROMIUM IONS	180
Aleksandar Jovanović, N. Knežević, N. Čutović, M. Đolić, N. Prlainović, Z. Veličković, M. Vuksanović IMPROVED TECHNOLOGY FOR PRODUCTION OF PE AND PP REGRANULATES	186
Jasmina Dedić, M. Mojsić, D. Lazarević, B. Stojčetočić, Ž. Šarkoćević A SHORT REVIEW OF TREPČA MINING WASTE IMPACT ON ENVIRONMENT	190
Senad Čeragić, H. Husić, V. Aganović INFLUENCE OF UNDERGROUND MINING WORKS OF THE OMAZIĆI BROWN COAL MINE BANOVIĆI ON THE SURFACE TERRAIN	195
Snežana Šerbula, J. Milosavljević, T. Kalinović, A. Radojević, J. Kalinović ARSENIC IN PARTICULATE MATTER ORIGINATED FROM MINING- METALLURGICAL PROCESSES	202
Nevena Ristić, M. Veličković, D. Voza THE ASSOCIATION BETWEEN SHORT-TERM EXPOSURE TO SO ₂ AND EMERGENCY ROOM ADMISSIONS IN URBAN AREA. CASE STUDY SERBIA	208
Miljan Marković, M. Gorgievski, N. Štrbac, K. Božinović, V. Grekulović, A. Mitovski, M. Zdravković ADSORPTION ISOTHERMS FOR COPPER IONS BIOSORPTION ONTO WALNUT SHELLS	214
Tanja Kalinović, J. Kalinović, S. Šerbula, J. Milosavljević, A. Radojević DETECTION OF THE TRAFFIC-RELATED POLLUTION BY THE ROADSIDE SOIL AND PLANT MATERIAL	219
Vladan Nedelkovski, S. Stanković, M. Radovanović, M. Antonijević SYNTHESIS AND CHARACTERISATION OF Ti/SnO ₂ -Sb-TYPE DSA ANODES FOR WASTEWATER TREATMENT	226
Maja Radić, M. Avdagić, B. Marković, K. Ademović, S. Avdagić, S. Avdagić PROTECTION AND PRESERVATION OF NATURAL RESOURCES	233
Mihajlo Stanković Annex 2: FAUNA OF MACROCRUSTACEA (INVERTEBRATA- ARTHROPODA) IN PERMANENT AND EPHEMERAL WATERS OF ZASAVICA WETLAND (SPECIAL NATURE RESERVES)	239

Mihajlo Stanković	OVERVIEW OF GALIS FAUNA (DIPTERA, HOMOPTER AND HYMENOPTER) – ZASAVICA RESERVE	246
Vanja Marković, M. Božanić, O. Arapović, V. Gojšina, A. Tatović, K. Stojanović	AQUATIC INSECTS DIVERSITY IN THE PART OF NERETVA RIVER DRAINAGE (BOSNIA AND HERZEGOVINA): PRELIMINARY RESULTS	253
Aleksandar Savić, I. Jelić, M. Šljivić-Ivanović, S. Dimović	RECYCLED CONCRETE AGGREGATE EFFECT ON SELF-COMPACTING CONCRETE AT LOW TEMPERATURES	258
Ivana Bjedov, D. Obratov-Petković, V. Stojanović, M. Nešić, D. Marisavljević	INVASIVE ALIEN SPECIES IN SERBIA: LEGISLATION, STRATEGY, AND PLANS	264
Jelena Majstorovic, M. Korac, S. Savkovic	THE EFFECTS OF GEOMECHANICAL RESEARCH “ <i>IN SITU</i> ” ON THE EXPLOITATION CONDITIONS OF THE OPEN PIT BAUXITE MINE “ZAGRAD”	271
Isidora Berezni, T. Marinković, V. Bežanović, M. Živančev, B. Batinić	ANALYSIS OF HOUSEHOLD'S E-WASTE AWARENESS, AND DISPOSAL BEHAVIOR IN NOVI SAD	276
Maja Radić, M. Avdagić, B. Markovic, K. Ademović, S. Avdagić, S. Avdagić	ECOLOGICAL ETHICS AND ECOLOGICAL EDUCATION	284
Vanja Marković, K. Zorić, M. Ilić, S. Đuretanović, M. Smederevac-Lalić	STONE CRAYFISH <i>AUSTROPOTAMOBIVUS TORRENTIUM</i> (Schrank, 1803) IN SOUTH-EASTERN SERBIA: CHILDREN KNOWLEDGE SURVEY	290
Ana Radojević, J. Milosavljević, T. Kalinović, J. Kalinović, S. Šerbula	THE IMPACT OF TEXTILE AND CLOTHES PRODUCTION ON THE ENVIRONMENT – PART I: ENVIRONMENTAL ISSUES	295
Ana Radojević, J. Milosavljević, T. Kalinović, J. Kalinović, S. Šerbula	THE IMPACT OF TEXTILE AND CLOTHES PRODUCTION ON THE ENVIRONMENT – PART II: WHAT CAN WE DO?	301
Giorgia Santini, V. Memoli, L. Santorufo, G. Di Natale, G. Maisto	EFFECTS OF UN-BIODEGRADABLE AND BIODEGRADABLE PLASTIC SHEETS ON ABIOTIC SOIL PROPERTIES	307
Milica Rajačić, D. Todorović, J. Krneta Nikolić, I. Vukanac, N. Sarap, M. Janković	RADIONUCLIDE CONTENT IN SAMPLES OF BERRIES	313
Ivana Mikavica, D. Randelović, J. Stojanović, J. Mutić	MICROPLASTIC OCCURRENCE IN URBAN AND SUBURBAN SOILS OF BOR, EASTERN SERBIA	319
Maja Nujkić, V. Stiklić, Ž. Tasić, S. Milić, D. Medić, A. Papludis, I. Đorđević	BIOSORPTION OF METAL IONS FROM SYNTHETIC SOLUTIONS USING DIFFERENT PARTS OF PLANT MATERIAL – A REVIEW	325
Nataša Đorđević, S. Mihajlović	ANALYSIS OF CHANGES OF MECHANICALLY ACTIVATED SODIUM CARBONATE SAMPLE	331
Vanja Trifunović, S. Milić, Lj. Avramović, R. Jonović, S. Đorđievski	ELECTRIC ARC FURNACE DUST – HAZARDOUS INDUSTRIAL WASTE WHOSE TREATMENT IS UNAVOIDABLE	336

Goran Milentijević, T. Marinković, M. Rančić, M. Milošević, I. Đuričković, A. D. Marinković, M. M. Milosavljević	OPTIMIZATION OF ZINC DIMETHYLDITHIOCARBAMATE SYNTHESIS PROCESS (ZIRAM)	343
Miloš Prokopijević, D. Spasojević, O. Prodanović, N. Pantić, D. Bartolić, K. Radotić, R. Prodanović	STABILITY OF SOYBEAN PEROXIDASE IMMOBILIZED ONTO HYDROGEL MICRO-BEADS FROM TYRAMINE-PECTIN	350
Nevena Pantić, M. Spasojević, M. Prokopijević, D. Spasojević, A. M. Balaž, R. Prodanović, O. Prodanović	COVALENT IMMOBILIZATION OF HORSERADISH PEROXIDASE ON NOVEL MACROPOROUS POLY(GMA-CO-EGDMA) FOR PHENOL REMOVAL	354
Dragica Spasojević, M. Prokopijević, O. Prodanović, N. Pantić, M. Stanković, K. Radotić, R. Prodanović	PREPARATION OF CROSSLINKED TYRAMINE-ALGINATE HYDROGEL USING EDC/NHS WITH SELF-IMMOBILIZED HRP	360
Jelena Mitrović, M. Radović Vučić, N. Velinov, S. Najdanović, M. Kostić, M. Petrović, A. Bojić	THE ROLE OF HYDROXYL AND SULFATE RADICALS IN THE UV ACTIVATED PERSULFATE DEGRADATION OF TEXTILE DYE RO16	364
Milica Petrović, T. Jovanović, S. Rančev, M. Radović Vučić, J. Mitrović, S. Najdanović, A. Bojić	ELECTROSYNTHESIZED CERIUM OXIDE CATALYST FOR ATMOSPHERIC PRESSURE PULSATING CORONA PLASMA DEGRADATION OF RB 5	369
Slobodan Najdanović, M. Petrović, M. Kostić, N. Velinov, J. Mitrović, D. Bojić, A. Bojić	PHOTOCATALYTIC DEGRADATION OF RANITIDINE BY BISMUTH OXO CITRATE	375
Ana Simonović, Ž. Tasić, M. Radovanović, M. Petrović Mihajlović, M. Antonijević	CAFFEINE AS A GREEN CORROSION INHIBITOR FOR COPPER IN SYNTHETIC BLOOD PLASMA SOLUTION	381
Dragana Medić, S. Milić, S. Alagić, M. Nujkić, A. Papludis, S. Đordjević, S. Dimitrijević	RECYCLING GOLD FROM WASTE PRINTED CIRCUIT BOARDS	387
Uroš Stamenković, S. Ivanov, I. Marković	CHARACTERIZATION OF CARBON AND LOW-ALLOY STEEL AFTER DIFFERENT HEAT TREATMENTS	393
Violeta Babić, M. Milenković, Z. Govedar, S. Stajić, B. Kanjevac	THE FOREST FIRES IN BULGARIA: THE TRENDS AND THE INFLUENCE OF TELECONNECTIONS	400
Martina Petković, M. Božović, A. Klikovac, D. Knežević	THE IMPORTANCE OF FIRE PROTECTION ON SHIPS	406
Darko Stojićević, Z. Živković, M. Saulić, T. Sekulić, V. Stupar	PESTICIDES – IMPACT ON HUMAN HEALTH AND THE ENVIRONMENT	413
Zlata Živković, M. Saulić, D. Stojićević, M. Jevtić Đorović, V. Stupar	ORGANIC AGRICULTURE: POTENTIAL OF THE FUTURE	419

Tatjana Anđelković, K. Kitanović, I. Kostić Kokić, B. Zlatković, D. Bogdanović CHANGES IN NITRATE AND NITRITE CONTENT IN FOR LETTUCE, CHARD AND SPINACH AFTER FREEZING	425
Ivana Perović, S. Brković, N. Zdošek, G. Tasić, M. Seović, S. Mitrović, J. Ciganović CAN LASER MODIFICATION OF COATED ELECTRODES IMPROVE THE HYDROGEN EVOLUTION REACTION IN ALKALINE ELECTROLYSERS?	430
Nikola Zdošek, S. Brković, I. Perović, M. Čurčić, S. Dimović, M. Vujković NEW GENERATION OF ELECTROCHEMICAL SUPERCAPACITORS	436
Nebojša Potkonjak, Đ. Čokeša, M. Marković THE HYDROGEN PRODUCTION ON Ni ELECTRODE CO-DEPOSITED WITH $\text{Co}+\text{V}_2\text{O}_5$: THE ELECTROCALYTIC SYNERGETIC EFFECT	441
Tatjana Ratknić, G. Šekularac, M. Ratknić, Z. Poduška, M. Aksić EFFECTS OF CLIMATE CHARACTERISTICS ON THE DIAMETER INCREMENT OF CEDAR IN THE CITY OF BELGRADE (SERBIA)	446
Natalija Čutović, M. Vuksanović, M. Milošević, M. Bugarčić, J. Bošnjaković, J. Gržetić, A. Marinković RECYCLED POLY(ETHYLENE TEREPHTHALATE) BASED- PLASTICIZER FOR PVC REGRANULATES PRODUCTION	452
Robert Vigi, G. Štrbac, D. Štrbac, M. Novaković SYNTHESIS OF NEW Ag DOPED CHALCOGENIDE GLASS FOR APPLICATION IN ENERGY CONVERSION AND MEMORY STORAGE	459
Tatjana Miljojčić, I. Jelić, M. Šljivić-Ivanović, S. Dimović, U. Ramadani SUSTAINABLE UTILIZATION OF CATHODE-RAY TUBE WASTE GLASS IN CEMENTITIOUS MATERIALS – A REVIEW	465
Dejan Riznić, A. Fedajev, A. Jevtić ECOTURISM AS A FORM OF GREEN ECONOMY	471
Vesna Đikanović, J. Vranković, K. Jovičić DIET OF TWO CYPRINID SPECIES, WHITE BREAM (BLICCA BJOERKNA) AND COMMON ROACH (RUTILUS RUTILUS) IN THE DANUBE RIVER, BELGRADE	477
Aleksandra Radić, D. Voza, Đ. Nikolić, M. Vuković ENVIRONMENTAL PERFORMANCE CLASSIFICATION OF BALKAN COUNTRIES BASED ON TOPSIS-SORT METHODOLOGY	483
Marijana Pantić, T. Maričić, S. Milijić CITIZEN PARTICIPATION OFFERS LESSONS TO CLIMATE CHANGE MANAGEMENT	489
Tanja Brdarić, D. Aćimović, B. Savić, N. Abazović, M. Čomor, M. Ječmenica Dučić, D. Maksin INVESTIGATION OF ZrO_2 AND $\text{ZrO}_2/\text{TiO}_2$ ELECTRODES BY CYCLIC VOLTAMMETRY	495
Branislava Savić, D. Aćimović, T. Brdarić, M. Ognjanović, D. Vasić Aničijević, M. Ječmenica Dučić, M. Simić TESTING THE ELECTROCHEMICAL BEHAVIOR OF BPA ON GC, WO_3 AND MWCNT ELECTRODES	499
Marijola Božović, M. Petković, S. Marković, B. Stojčetović DISASTER RISK MANAGEMENT FRAMEWORK	504
Author Index	510

ELECTRIC ARC FURNACE DUST – HAZARDOUS INDUSTRIAL WASTE WHOSE TREATMENT IS UNAVOIDABLE

Vanja Trifunović^{1*}, Snežana Milić², Ljiljana Avramović¹, Radojka Jonović¹,
Stefan Đordjević¹

¹Mining and Metallurgy Institute Bor, Zeleni bulevar 35, 19210 Bor, SERBIA

²University of Belgrade, Technical Faculty in Bor, V.J. 12, 19210 Bor, SERBIA

*vanja.trifunovic@irmbor.co.rs

Abstract

Electric arc furnace dust (EAF dust), which is formed as a by-product of the steel production process from secondary raw materials, contains significant amounts of Zn, Fe, Pb, Cd, Ca, K, Cr, Mn, Si, Mg, Na, Ni, Cu, F, Cl, etc. Due to the content of heavy metals in the EAF dust, which in its inadequate disposal and the impact of atmospheric conditions can lead to a negative impact on the environment, it is characterized in the world as hazardous industrial solid waste. For environmental protection, it is necessary to do a treatment of the EAF dust so that it can be safely disposed of in a landfill without negative effects on the environment and human health. In addition to environmental benefits, the EAF dust treatments can also have economic benefits.

Keywords: EAF dust, hazardous waste, environmental protection

INTRODUCTION

World production of crude steel takes place in two ways: processing of primary raw materials, primarily iron oxide ore, melting in the Basic Oxygen Furnace (BOF) and processing of iron-based secondary raw materials by melting in an Electric Arc Furnace (EAF) [1–4].

Using secondary raw materials, i.e. scrap iron, as a raw material for steel production, and electricity as an energy source, steel production in electric arc furnaces has become more prominent than any other steel production process in the world [5–11]. The main sources of secondary raw materials for steel production are construction material waste, old cars, appliances and household waste, which means that waste can contain a large number of metals, plastics and rubber, glass, paint, oil, and even salt [2,5–8,12]. Figure 1 shows an example of scrap iron used as a batch for an electric arc furnace [13].

During the melting process of scrap iron in the electric arc furnace at a temperature of 1600°C, a by-product of the process-dust from the electric arc furnace is forming. Emission of the electric arc furnace dust (EAF dust) is proportional to the steel production so that with the increasing steel production there is an increase in the amount of the EAF dust: for each ton of produced crude steel about 10–20 kg of the electric arc furnace dust is also produced. About 5–7 million tonnes of dust are generated annually in the world, of which 0.5–0.9 million tonnes of dust come from steel plants in Europe [2,4–6,8,10,11,14,15].



Figure 1 Scrap iron used as a batch in the steel production process from secondary raw materials [13]

Formation of the EAF dust occurs since volatile elements, such as Zn, Pb and Cd in the batch of the furnace mostly evaporate in the early stages of the operation, and eventually, all of them go into the gas phase. Other elements, such as Fe, Cr, Ni and Mn, are generated from gaseous products during the refining period, mainly due to oxygen injection. Volatile components are rapidly converted into the corresponding oxides in contact with air after the combustion and cooling of the exhaust gases [1,2,5,7]. Oxides in the slag mixed with metal vapors and generated dust are extracted from the furnace by hot gas flow, to a gas purification system where gases are cooled down, and finally, the EAF dust is formed which can be collected in a gravity collector, electrostatic dust collector, or filter bags [1,5,14].

EAF DUST CHARACTERIZATION

Electric arc furnace dust occurs in the form of very fine reddish-brown or dark brown particles, which can spread in the air [5,7]. Due to the difference in the procedures for obtaining steel in an electric arc furnace and the methods of collecting the EAF dust, the physical characteristics of EAF dust will vary within the appropriate range. For example, depending on the way the EAF dust is collected, the size of its particles will vary. When the EAF dust is collected with a gravity collector, 85% of dust particles are $<10\ \mu\text{m}$, by collecting in filter bags, 90% of particles are $<50\ \mu\text{m}$, and in an electrostatic collector, more than 90% of dust particles are $<100\ \mu\text{m}$ in size [5].

To a large extent, the composition of the EAF dust depends on the operating conditions of the electric arc furnace, the characteristics of the scrap iron batch, the working period, the specifications of the produced steel, and is also specific to each plant [1,2,6,8,14]. The composition of EAF dust can be variable and can change from day to day in the same steel plant [1]. EAF dust is the final result of a series of physical and chemical phenomena through which substances that create EAF dust pass and they define its physical aspect, chemical and mineral composition. Basically, EAF dust should consist of iron oxides, however, due to the presence of different types of scrap iron, which contains several different elements, its composition becomes complex [1,11]. It can be seen from Table 1 that the chemical composition of EAF dust from different sources can vary significantly.

Table 1 Comparative chemical composition of the EAF dust from different sources

Elements (%)													Ref.
Zn	Fe	Pb	Ca	Cd	Na	K	Cl ⁻	Mn	Si	Cr	Cu	Al	
22.73	14.40	4.22	13.32	0.09	1.25	1.61	4.75	1.00	1.45	-	-	-	[3]
29.10	24.00	3.64	3.16	-	-	1.76	1.56	4.11	1.57	-	0.25	0.52	[12]
17.80	4.20	2.34	0.80	-	-	-	1.60	0.62	8.14	-	-	0.73	[17]
33.00	26.50	2.17	0.90	-	-	-	-	2.30	-	0.20	0.20	-	[19]
24.24	34.38	1.76	2.27	-	-	-	4.72	-	-	-	-	-	[14]
7.79	37.29	1.16	5.33	-	2.78	3.11	2.63	1.18	2.21	0.15	-	-	[18]

As for the mineral composition, because the EAF dust is formed in oxidative conditions, most metals are present in the form of oxides. Zinc occurs in the form of ZnO and ZnFe₂O₄, while iron mainly occurs in the oxide form as Fe₃O₄ and Fe₂O₃ [1,2,6,8,11]. Calcium is present in the form of CaO, CaCO₃, calcium silicate, and calcium aluminosilicate. Lead, copper, chromium, and nickel can occur in the form of oxides, chlorides, sulfides, or sulfates. In recent years, there has been an increase in chlorine levels in the EAF dust due to an increase in impurities containing chlorine in scrap iron (most commonly rubber, paints, polymers, etc.), and it occurs in the form of alkali metal chlorides [1,2].

EFFECT OF EAF DUST ON THE ENVIRONMENT

The presence of metals such as Pb, Zn and Cd in the EAF dust can pose a threat to the environment and human health due to the mobility of these toxic elements, and for this reason, EAF dust is considered hazardous industrial solid waste in many countries [1,4–6,8]. According to the Environmental Protection Agency (EPA) in the United States it is characterized as hazardous solid industrial waste K061 [4], according to the Brazilian standard ABNT 10004: 2004, EAF dust is listed as hazardous waste from a specific source K061 [5,20], in the catalog of the European Union Waste [21] it is included in the list of hazardous substances with the code 10 02 13, and according to the European catalog of waste [22] it is classified as a hazardous waste of category 10 02 07 [6]. It is also considered an air pollutant because the gas purification system captures only 70% of the EAF dust, while the rest goes into the atmosphere. Therefore, EAF dust is a solid hazardous waste that has the potential for long-term environmental pollution [23].

The pollution potential of this type of waste lies in the possibility of self-leaching elements such as Zn, Cu, Cr, Ni, Pb, Cd, F, and Cl. Factors that play a significant role in increasing the solubility of heavy metals from the EAF dust in the environment are pH, redox potential, and temperature. Controlling these factors, the risk of environmental contamination with heavy metals from the EAF dust can be reduced [23]. Disposal of this hazardous waste has become a serious problem in recent years since increasing steel production increases the amount of hazardous waste that needs to be adequately treated, in prescribed landfills, to prevent its negative impact on the environment and human health. It is understood that when disposing of the EAF dust in hazardous waste landfills, the EAF dust must be protected from rain to prevent the formation of leachate that could contaminate the land and watercourses in the surrounding area [4,24]. Figure 2 shows the “mountain” of disposed EAF dust generated from

steel production in Egypt [14]. As can be seen from the figure, this method of disposal is not adequate since hazardous waste is completely exposed to external atmospheric influences and it is very easy to contaminate the environment with heavy metals from the dust [4,15,24].




Figure 2 a) “Mountain” of the EAF dust generated from steel production in Egypt; b) a representative sample of the EAF dust [25]

EAF DUST TREATMENT TO ENVIRONMENTAL PROTECTION

For environmental protection and the protection of human health, it is necessary to do a treatment of this type of hazardous waste. Certain treatments of the EAF dust, in addition to reducing the amount of hazardous waste and/or its transformation into non-hazardous waste to protect the environment, would also make savings on the cost of its disposal. Also, in addition to environmental benefits, economic benefits could be achieved by extracting zinc for commercial quality products and using iron concentrates suitable as raw materials in steel or pig iron production [1,6,14].

Although very rich in iron, EAF dust cannot be recycled directly into the steel production process, because it contains large amounts of Zn (from 2–40% [3,8,12,14,17–19]), which negatively affects the electric arc furnace and thus endangers the economy of the furnace. Although chrome and lead are not problematic for the steel production process itself, they are toxic to humans and their content is regulated by the European Commission. Consequently, metal separation has become a desirable option, as the EAF dust is a quality secondary raw material for zinc production – EAF dust contains approximately 7% of world zinc production that would otherwise be thrown in landfills. With the constant increase in demand for zinc and with limited possibilities of its substitution in materials and the depletion of reserves from its primary sources, the need for secondary sources of zinc is growing. Therefore, the EAF dust treatment process must be able to remove Zn from the dust, while the remaining valuable metals are compatible with iron and steel so that the iron-containing material can be recovered and recycled in the steel production process, or to obtain a solid residue that can be safely disposed of as non-hazardous waste [8]. Efficient extraction of zinc from secondary sources has several advantages such as: saving intact resources and fossil resources used to supply energy in primary mining processes, increased resource efficiency, reduced landfill disposal and loss of zinc or any metal returned to landfill, waste remediation, as well as reduction of environmental impact [23].

Given the potential environmental and economic benefits from the EAF dust, it is important to develop appropriate processes for its treatment. The chemical composition of the EAF dust, as well as its available amount, provides opportunities and conditions for deciding on the best sustainable method for its treatment. For the recycling of the EAF dust for zinc separation to be economically viable, the dust must contain Zn concentrations higher than 15% [20]. In general, there are two basic processes in the EAF dust treatment: the pyrometallurgical process and the hydrometallurgical process. Both processes have their advantages and disadvantages. Pyrometallurgical processes require high temperatures, expensive equipment, and large capacity plants, which leads to high energy consumption. For the treatment of the EAF dust by pyrometallurgical processes, the minimum zinc content in the dust must be >15% to ensure the economy of the process. By applying this process, a product ZnO of low commercial value is obtained, which requires additional hydrometallurgical treatment. Also, in this process other solid residues are formed and need additional treatment before disposal [2,5]. On the other hand, the hydrometallurgical process can be economical with a zinc content in dust lower than 15%. These processes have significantly lower capital and operating costs compared to pyrometallurgical ones, can be installed in situ, and commercial quality products can be obtained. However, large amounts of wastewaters are generated that require further treatment to be safely discharged to recipients. Solid residues of these processes may or may not be subsequently treated before disposal. Extraction of zinc from the EAF dust by the hydrometallurgical process is very complicated and consists of several related processes, such as leaching, purification of pregnant leaching solution, and separation of metals/compounds [3,5,6,8,14].

Other methods of recycling the EAF dust were also investigated, not to separate valuable metals and obtain non-hazardous waste, but for direct disposal of the dust by incorporating dust containing zinc in a concentration of less than 15% in existing products. Products in which EAF dust can be incorporated are blue shales [15] for use in the cement industry, bricks [4], kaolin clay used for 3D printing in the construction industry [10], ceramic tiles, etc. In addition to saving on the costs of disposing of the EAF dust, its incorporation into existing products leads to improved physical characteristics of these products [4,10,15].

CONCLUSION

Dust from the electric arc furnace generated in the steel production process from secondary raw materials contains significant amounts of Zn, Fe, and Pb, as well as variable amounts of Cd, Ca, Cr, Mn, Si, Mg, Na, F, Cl, etc. Due to the content of heavy metals in the EAF dust, which can contaminate the surrounding lands and watercourses due to its inadequate disposal, and the influence of atmospheric conditions, it is characterized in the world as hazardous industrial solid waste.

EAF dust treatment processes for the recovery of valuable metals can be pyrometallurgical and hydrometallurgical processes, depending on dust chemical composition and the amount available for processing. All processes have their advantages and disadvantages.

For environmental protection, it is necessary to do the treatment of the EAF dust so that it can be safely disposed of in a landfill without negative effects on the environment and human

health. In addition to environmental benefits, the EAF dust treatment can also have economic benefits.

The impact of incorporated EAF dust on the environment has not been sufficiently investigated.

ACKNOWLEDGMENTS

The authors are grateful to the Ministry of Education, Science and Technological development of the Republic of Serbia for financial support (Grant No. 451-03-68/2022-14/200052 and Grant No. 451-03-68/2022-14/200131).

REFERENCES

- [1] P. Keglevich de Buzina, N. Heckb, A. Vilela, J. Mater. Res. Technol; 6 (2) (2017) 194–202.
- [2] C. Pickles, Miner. Eng; 22 (2009) 977–985.
- [3] V. Montenegro, S. Agatzini-Leonardou, P. Oustadakis, *et al.*, Waste Biomass Valori; 7 (6) (2016) 1531–1548.
- [4] E. Dominguez, R. Ullmann, Appl. Clay Sci; 6 (1996) 237–249.
- [5] J. Wang, Y. Zhang, K. Cui, *et al.*, J. Cleaner Prod; 298 (2021) 126788.
- [6] F. Kukurugya, T. Vindt, T. Havlík, Hydromet; 154 (2015) 20–32.
- [7] V. Silva, J. Silva, *et al.*, J. Mater. Res. Technol; 8 (6) (2019) 5504–5514.
- [8] P. Hazaveh, S. Karimia, *et al.*, Ecotoxicol. Environ. Saf; 202 (2020) 110893.
- [9] O. Ruiz, C. Clemente, M. Alonso, *et al.*, J. Hazard. Mater; 141 (2007) 33–36.
- [10] E. Ordoñez, S. Neves Monteiro, H. Colorado, Mater. Des; 217 (2022) 110617.
- [11] V. Trifunović, Lj. Avramović, R. Jonović, *et al.*, Proceedings of the 52nd International October Conference on Mining and Metallurgy, November 29th–30th, Bor, Serbia (2021) 209–212.
- [12] C. Lanzerstorfer, J. Cleaner Prod; 174 (2018) 1–6.
- [13] M. Siame, J. Kaoma, N. Hlabangana, *et al.*, S. Afr. J. Chem. Eng; 27 (2019) 35–42.
- [14] R. Fujimoto, K. Maruyama, T. Miki, *et al.*, Hydromet; 159 (2016) 120–125.
- [15] S. Cholake, R. Farzana, T. Numata, *et al.*, J. Cleaner Prod; 171 (2018) 1128–1139.
- [16] P. Oustadakis, P. Tsakiridis, A. Katsiapi, *et al.*, J. Hazard. Mater; 179 (2010) 1–7.
- [17] P. Dvořák, H.N. Vu, J. Pol. Min. Eng. Soc; (2017) 195–199.
- [18] H. Wang, J. Gao, W. Liu, *et al.*, Hydrom; 166 (2016) 1–8.
- [19] T. Havlík, B. Vidor e Souza, *et al.*, J. Hazard. Mater; B135 (2006) 311–318.
- [20] J. Araújo, V. Schalch, J. Mater. Res. Technol; 3 (3) (2014) 274–279.
- [21] Review of the European List of Waste
https://ec.europa.eu/environment/pdf/waste/low_review_oekopol.pdf
- [22] European Catalog of Waste, Available on the following link:

<https://www.scribd.com/doc/26038172/Evropski-katalog-otpada>

- [23] M. Al-harashseh, J. Al-Nuairat1, A. Al-Otoom, *et al.*, J. Environ. Chem. Eng; 7 (2019) 102856.
- [24] E. Ledesma, A. Lunar, J. Ayuso, *et al.*, Constr. Build. Mater; 183 (2018) 365–375.
- [25] R.M. Khattab, M.M. El-Sayed Seleman, M.F. Zawrah, Ceram. Int; 43 (2017) 12939–12947.

tfbor.bg.ac.rs
eco.tfbor.bg.ac.rs

EcoTER'22



ISBN 978-86-6305-123-2